

ON KÖNIG-EGERVÁRY SQUARE-STABLE GRAPHS

VADIM E. LEVIT AND EUGEN MANDRESCU

ABSTRACT. The *stability number* of a graph G , denoted by $\alpha(G)$, is the cardinality of a maximum stable set, and $\mu(G)$ is the cardinality of a maximum matching in G . If $\alpha(G) + \mu(G)$ equals its order, then G is a König-Egerváry graph.

In this paper we deal with *square-stable graphs*, i.e., the graphs G enjoying the equality $\alpha(G) = \alpha(G^2)$, where G^2 denotes the second power of G . In particular, we show that a König-Egerváry graph is square-stable if and only if it has a perfect matching consisting of pendant edges, and in consequence, we deduce that well-covered trees are exactly the square-stable trees.

Keywords: stable set, square-stable graph, well-covered graph, matching.

2010 *Mathematics Subject Classification:* 05C69, 05C76.

1. INTRODUCTION

All the graphs considered in this paper are simple, i.e., are finite, undirected, loopless and without multiple edges. For such a graph $G = (V, E)$ we denote its vertex set by $V = V(G)$ and its edge set by $E = E(G)$. If $X \subset V$, then $G[X]$ is the subgraph of G spanned by X . By $G - W$ we mean the subgraph $G[V - W]$, if $W \subset V(G)$.

A set of pairwise non-adjacent vertices is a *stable set* of G . The *stability number* of G , denoted by $\alpha(G)$, is the size of a maximum stable set in G .

Let $\Omega(G)$ stand for the set $\{S : S \text{ is a maximum stable set of } G\}$ and $\text{core}(G) = \cap\{S : S \in \Omega(G)\}$ [11].

$\theta(G)$ is the *clique covering number* of G , i.e., the minimum number of cliques whose union covers $V(G)$.

Recall that

$$i(G) = \min\{|S| : S \text{ is a maximal stable set in } G\},$$

and

$$\gamma(G) = \min\{|D| : D \text{ is a minimal domination set in } G\}.$$

A *matching* is a set of non-incident edges of G ; a matching of maximum cardinality $\mu(G)$ is a *maximum matching*, and a *perfect matching* is a matching covering all the vertices of G .

G is called a *König-Egerváry graph* provided $\alpha(G) + \mu(G) = |V(G)|$ [3, 24]. Various properties of König-Egerváry graphs are presented in [8, 9, 12, 14, 15, 16, 17, 18, 19].

According to a well-known result of König [7], and Egerváry [4], any bipartite graph is a König-Egerváry graph. This class includes non-bipartite graphs as well (see, for instance, the graphs H_1 and H_2 in Figure 1).




Figure 1: Only H_3 is not a König-Egerváry graph.

It is easy to see that if G is a König-Egerváry graph, then $\alpha(G) \geq \mu(G)$, and that a graph G having a perfect matching is a König-Egerváry graph if and only if $\alpha(G) = \mu(G)$.

The neighborhood of a vertex $v \in V$ is the set

$$N(v) = \{w : w \in V, vw \in E\}, N[v] = N(v) \cup \{v\},$$

and $N(A) = \cup\{N(v) : v \in A\}$, for $A \subset V$.

If $G[N(v)]$ is a complete subgraph in G , then v is a *simplicial vertex* of G , and by $\text{simp}(G)$ we mean the set of all simplicial vertices of G . A maximal clique in G is called a *simplex* if it contains at least a simplicial vertex of G [2]. G is said to be *simplicial* if every vertex of G is simplicial or it is adjacent to a simplicial vertex of G [2]. If $|N(v)| = |\{w\}| = 1$, then v is a *pendant vertex* and vw is a *pendant edge* of G .

By K_n and P_n we denote the complete graph and the cordless path on $n \geq 1$ vertices, respectively. $K_{n,m}$ is the complete bipartite graph with two maximal stable sets of cardinalities n and m .

G is *well-covered* [20] if every maximal stable set of G is also a maximum stable set, i.e., it is in $\Omega(G)$. G is called *very well-covered* [5] provided G is well-covered without isolated vertices, and $|V(G)| = 2\alpha(G)$.

The following results will be used in the sequel.

Proposition 1 *If G is a graph of order $n \geq 2$, then:*

1. [10] G is very well-covered if and only if it is a well-covered König-Egerváry graph;
2. [10] a connected König-Egerváry graph is well-covered if and only if it is very well-covered.
3. [1] $H = G - N[v]$ is well-covered and $\alpha(H) = \alpha(G) - 1$, for every $v \in V(G)$, whenever G is a non-complete well-covered graph.

The distance between two vertices v, w of a graph G is denoted by $\text{dist}_G(v, w)$, or $\text{dist}(v, w)$ if no ambiguity. G^2 denotes the *second power* of G , i.e., the graph with

$$V(G^2) = V(G), \quad E(G^2) = E(G) \cup \{uv : u, v \in V(G), \text{dist}_G(u, v) = 2\}.$$

Clearly, any stable set of G^2 is stable in G , as well, while the converse is not generally true. Therefore, we may assert that $1 \leq \alpha(G^2) \leq \alpha(G)$. Notice that the both bounds are sharp. For instance, if:

- $G = K_{1,n}$, and $n \geq 2$, then $\alpha(G) = n > \alpha(G^2) = 1$;
- $G = P_4$, then $\alpha(G) = \alpha(G^2) = 2$.




Figure 2: G_1 and G_2 are square-stable graphs, while G_3 is not square-stable.

In this paper we characterize König-Egerváry graphs satisfying the equality $\alpha(G) = \alpha(G^2)$. The graphs enjoying this property are called *square-stable* [13]. The investigation of these graphs was started in [22] and continued in [13].

2. RESULTS

It is evident that G and G^2 are simultaneously connected or disconnected. In addition, if $H_i, 1 \leq i \leq k$, are the connected components of graph G , then $S \in \Omega(G)$ if and only if $S \cap V(H_i) \in \Omega(H_i), 1 \leq i \leq k$. Hence, we may conclude that a disconnected graph is square-stable if and only if each of its connected components is square-stable. Therefore, in the rest of the paper all the graphs are connected, unless otherwise stated.

In general, $\alpha(G^2) \leq \theta(G^2) \leq \gamma(G) \leq i(G) \leq \alpha(G) \leq \theta(G)$.

Theorem 2 *For a graph G the following statements are equivalent:*

- (i) *every vertex of G belongs to exactly one simplex of G ;*
- (ii) *G satisfies $\alpha(G) = \alpha(G^2)$, i.e., G is square-stable;*
- (iii) *G satisfies $\theta(G) = \theta(G^2)$;*
- (iv) *G satisfies $\alpha(G^2) = \theta(G^2) = \gamma(G) = i(G) = \alpha(G) = \theta(G)$;*
- (v) *G is simplicial and well-covered;*
- (vi) *there exists some $S \in \Omega(G)$ such that $\text{dist}_G(a, b) \geq 3$ holds for all distinct $a, b \in S$.*

Recall that: the equivalence of the assertions (i), (ii), (iii), (iv), was proved in [22], the fact that "(i) $\iff$ (v)" was shown in [21], while "(ii) $\iff$ (vi)" was done in [13].

For square-stable graphs there are interesting connections between properties of vertices in G and G^2 correspondingly.

Proposition 3 *If G is a square-stable graph and $v \in V(G)$, then the following assertions are true:*

- (i) *v is in some $S \in \Omega(G^2)$ if and only if v is a simplicial vertex in G ;*
- (ii) *v belongs to all maximum stable sets of G^2 if and only if v is the unique simplicial vertex in its simplex in G .*

Proof. (i) Let v belong to some $S \in \Omega(G^2)$. If v is not simplicial in G , then there are $x, y \in N_G(v)$, such that $xy \notin E(G)$. In addition,

$$N_G(x) \cap S = \{v\} = N_G(y) \cap S,$$

because $\text{dist}_G(v, u) \geq 3$, for each $u \in S - \{v\}$. Since S belongs also to $\Omega(G)$, we infer that $(S \cup \{x, y\}) - \{v\}$ must be stable in G and of cardinality greater than $|S|$, thus contradicting $|S| = \alpha(G)$.

Conversely, let v be a simplicial vertex in G . Suppose that $S \in \Omega(G^2)$, and let $\{u\} = S \cap N_G[v]$. Hence, $\text{dist}_G(x, v) \geq 3$ for all $x \in S - \{u\}$. Otherwise, there exists a vertex $x_0 \in S - \{u\}$ such that $\text{dist}_G(v, x_0) = 2$, which implies $\text{dist}_G(u, x_0) = 2$, because v is simplicial in G . Therefore,

$$S' = (S - \{u\}) \cup \{v\} \in \Omega(G^2) \text{ and } v \in S'.$$

(ii) Suppose that a vertex v belongs to all maximum stable sets of G^2 . According to (i), v must be a simplicial vertex in G . If some $u \in N_G(v)$ is also simplicial in G , then again by (i), u belongs to some $S \in \Omega(G^2)$. Clearly, $v \notin S$ and this contradicts the hypothesis on v .

Conversely, let v be the unique simplicial vertex in its simplex. Assume that there is some $S \in \Omega(G^2)$ such that $v \notin S$. Since G is square-stable, $S \in \Omega(G)$. Clearly, there exists $u \in S \cap N_G(v)$, because, otherwise, $S \cup \{v\}$ is stable in G and $|S \cup \{v\}| > \alpha(G)$. Hence, $S \cap N_G(u) = \emptyset$, and each $x \in S$ satisfies $\text{dist}_G(x, u) \geq 2$.

The vertex u is not simplicial, since v is the unique simplicial vertex of its simplex and $u \in N_G(v)$. Consequently, there is $y \in N_G(u) - N_G[v]$.

Now we get that $(S \cup \{v, y\}) - \{u\}$ is stable in G , because $S \in \Omega(G^2)$ means that $\text{dist}_G(x, u) \geq 3$, which implies $\text{dist}_G(x, y) \geq 2$ for all $x \in S - \{u\}$. Thus, we get that

$$|(S \cup \{v, y\}) - \{u\}| > \alpha(G),$$

in contradiction with the definition of $\alpha(G)$. In summary, v must belong to all maximum stable sets of G^2 . ■

The graph in Figure 3 illustrates interconnections between $\text{simp}(G)$ and $\text{core}(G^2)$ implied by Proposition 3.




Figure 3: G is square-stable and $v_2, v_4, v_5, v_8 \in \text{simp}(G)$, but only $v_2, v_5 \in \text{core}(G^2)$.

Clearly, any complete graph is square-stable. Moreover, since $K_n^2 = K_n$, we get that

$$|\Omega(K_n)| = |\Omega(K_n^2)| = |\{\{v\} : v \in V(K_n)\}| = n.$$

Let us notice that the equality $|\Omega(G^2)| = 1$ does not ensure that G is square-stable, e.g., the graph G from Figure 4 is not square-stable, but $|\Omega(G^2)| = 1$.




Figure 4: $\alpha(G) = |\{x, y, z\}| = 3$, while $\Omega(G^2) = \{\{x, y\}\}$.

In general, a graph having a unique perfect matching is not necessarily square-stable. For instance, $K_3 + e$ has a unique perfect matching, but is not square-stable, because $(K_3 + e)^2 = K_4$.

Further, we pay attention to graphs having a perfect matching consisting of pendant edges, which is obviously unique.

Lemma 4 *If G has a perfect matching consisting of pendant edges, then G is square-stable. Moreover, $\Omega(G^2) = \{S_0\}$, where $S_0 = \{v : v \text{ is pendant in } G\}$.*

Proof. Clearly, the set S_0 is stable in G , and because

$$|V(G)|/2 = \mu(G) = |S_0| \leq \alpha(G) \leq |V(G)| - \mu(G) = |V(G)|/2,$$

we infer that $S_0 \in \Omega(G)$. Since $\text{dist}_G(a, b) \geq 3$ holds for any $a, b \in S_0$, we get that $S_0 \in \Omega(G^2)$, i.e., G is square-stable, by Theorem 2(vi).

Since a pendant vertex is the unique simplicial vertex in its simplex, Proposition 3 implies that S_0 is included in all maximum stable sets of G^2 . So, we may conclude that $\Omega(G^2) = \{S_0\}$, because $S_0 \in \Omega(G^2)$. ■

Notice that there are square-stable graphs with more than one maximum stable set, and having no perfect matching; e.g., the graph in Figure 5.




Figure 5: A square-stable graph G and its G^2 . G is not very well-covered.

Theorem 5 *For a König-Egerváry graph G on $n \geq 2$ vertices the following assertions are equivalent:*

- (i) G square-stable;
- (ii) G has a perfect matching consisting of pendant edges;
- (iii) G is very well-covered with exactly $\alpha(G)$ pendant vertices.

Proof. (i) $\Rightarrow$ (ii) By Theorem 2(v), G is well-covered, and according to Proposition 1(2), it is also very well-covered. Hence, we obtain that $\alpha(G) = \mu(G) = n/2$, and G has a perfect matching M . Let

$$S_0 = \{a_i : 1 \leq i \leq \alpha(G)\} \in \Omega(G^2)$$

and $b_i \in V(G) - S_0$ be such that $a_i b_i \in M$ for $1 \leq i \leq \alpha(G)$. The inequality $\text{dist}_G(v, w) \geq 3$ holds for any $v, w \in S_0$, because otherwise $vw \in E(G^2)$, and this contradicts the stability of S_0 in G^2 .

If $b_j \in N_G(a_i)$ for some $i \neq j$, it follows that $\text{dist}_G(a_i, a_j) = 2$, in contradiction with $\text{dist}_G(a_i, a_j) \geq 3$. Therefore, $N_G(a_i) = \{b_i\}$, $1 \leq i \leq \alpha(G)$, i.e., M consists of only pendant edges.

(ii) $\Rightarrow$ (iii) By Lemma 4, it follows that G is square-stable. According to Theorem 2(v), G is well-covered, and Proposition 1(2) finally assures that G is very well-covered.

(iii) $\Rightarrow$ (i) Since G is very well-covered with exactly $\alpha(G)$ pendant vertices, we infer that

$$S_0 = \{v : v \text{ is a pendant vertex in } G\} \in \Omega(G)$$

and also that the matching $M = \{vw : vw \in E(G), v \in S_0\}$ is perfect and consists of only pendant edges. According to Lemma 4, it follows that G is square-stable. ■

Let us remark that:

- well-covered König-Egerváry graphs may not be square-stable, e.g., C_4 ;
- a König-Egerváry graph with a unique perfect matching is not always square-stable, e.g., P_6 (by the way, it is also a tree) and $K_3 + e$;
- a non-König-Egerváry graph with a unique perfect matching M may be square-stable, even if M does not consist of only pendant edges (for instance, see the graph in Figure 6).




Figure 6: G is square-stable and has a unique perfect matching containing not only pendant edges.

Theorem 5 is true for bipartite graphs as well, since any bipartite graph is also a König-Egerváry graph. For trees, Theorem 5 leads to an extension of the characterization that Ravindra gave to well-covered trees in [23].

Corollary 6 *If T is a tree of order $n \geq 2$, then the following statements are equivalent:*

- (i) T is well-covered;
- (ii) T is very well-covered;
- (iii) T has a perfect matching consisting of pendant edges;
- (iv) T is square-stable.

Let us notice that the equivalences appearing in Corollary 6 fail for bipartite graphs. For instance, the graph in Figure 7 is very well-covered, but is not square-stable.




Figure 7: A very well-covered bipartite graph, which is not square-stable.

Lemma 7 *If G is square-stable with 2 vertices at least, then $\alpha(G) \leq \mu(G)$.*

Proof. By Theorem 2(vi) there exists a maximum stable set

$$S = \{v_i : 1 \leq i \leq \alpha(G)\}$$

in G such that $\text{dist}_G(a, b) \geq 3$ for all $a, b \in S$. It means that for every $i \in \{1, 2, \dots, \alpha(G) - 1\}$ there is a shortest path of length 3 at least from v_i to $v_{\alpha(G)}$: $v_i, w_i, \dots, w^i, v_{\alpha(G)}$. All the vertices $w_i, 1 \leq i \leq \alpha(G) - 1$ and w^1 are pairwise distinct, otherwise there will be a pair of vertices in S at distance 2. Hence

$$M = \{v_i w_i : 1 \leq i \leq \alpha(G) - 1\} \cup \{v_{\alpha(G)} w^1\}$$

is a matching in G . Thus $\alpha(G) = |S| \leq |M| \leq \mu(G)$. ■

Proposition 8 *Let G^2 be a König-Egerváry graph with 2 vertices at least. Then G is square-stable if and only if G is a König-Egerváry graph with a perfect matching.*

Proof. The following inequalities are true for every graph G :

$$\mu(G) \leq \mu(G^2) \text{ and } \alpha(G^2) \leq \alpha(G).$$

Since G^2 is a König-Egerváry graph, $\mu(G^2) \leq \alpha(G^2)$. Consequently,

$$\mu(G) \leq \mu(G^2) \leq \alpha(G^2) \leq \alpha(G).$$

If G is square-stable, then these inequalities together with Lemma 7 give

$$\mu(G) = \mu(G^2) = \alpha(G^2) = \alpha(G).$$

Moreover,

$$|V(G)| = \mu(G^2) + \alpha(G^2) = \mu(G) + \alpha(G),$$

which means that G is a König-Egerváry graph. In addition, G has a perfect matching, because $\mu(G) = \alpha(G)$.

Conversely, if G is a König-Egerváry graph with a perfect matching, then

$$\mu(G) + \alpha(G) = |V(G)| = \mu(G^2) + \alpha(G^2) \text{ and } \mu(G) = \mu(G^2).$$

Thus $\alpha(G) = \alpha(G^2)$, i.e., G is a square-stable graph. ■

It is worth noticing that if G is square-stable, then it is not enough to know that $\mu(G) = \alpha(G)$ in order to be sure that G is a König-Egerváry graph. For instance, see Figure 8.




Figure 8: G is a square-stable non-König-Egerváry graph: $\mu(G) = \alpha(G) = 2$.

Combining Propositions 1, 8 and Theorem 5, we obtain the following.

Corollary 9 (i) *G is square-stable and very well-covered if and only if G is a König-Egerváry graph with exactly $\alpha(G)$ pendant vertices.*

(ii) *If G is square-stable, then either G and G^2 are König-Egerváry graphs or both of them are not.*

Another consequence of Theorem 5 is the following extension of the characterization that Finbow *et al.* gave in [6] for well-covered graphs having the girth ≥ 6 .

Theorem 10 *Let G be a graph of girth ≥ 6 , which is isomorphic to neither C_7 nor K_1 . Then the following assertions are equivalent:*

- (i) G is well-covered;
- (ii) G has a perfect matching consisting of pendant edges;
- (iii) G is very well-covered;
- (iv) G is a König-Egerváry graph with exactly $\alpha(G)$ pendant vertices, and $\text{core}(G) = \emptyset$;
- (v) G is a König-Egerváry square-stable graph.

Proof. The equivalences $(i) \Leftrightarrow (ii) \Leftrightarrow (iii)$ are proved in [6]. In [10] it is shown that $(iii) \Leftrightarrow (iv)$. Finally, $(ii) \Leftrightarrow (v)$ is true according to Lemma 4 and Theorem 5. ■




Figure 9: Square-stable König-Egerváry graphs.

3. CONCLUSIONS

In this paper we concentrated on König-Egerváry graphs enjoying the equality $\alpha(G^2) = \alpha(G)$. It seems to be interesting to study other families of graphs satisfying the same property.

References

- [1] S. R. Campbell, M. D. Plummer, *On well-covered 3-polytopes*, Ars Combinatoria **25-A** (1988) 215-242.
- [2] G. H. Cheston, E. O. Hare, R. C. Laskar, *Simplicial graphs*, Congressus Numerantium **67** (1988) 105-113.
- [3] R. W. Deming, *Independence numbers of graphs - an extension of the König-Egerváry theorem*, Discrete Mathematics **27** (1979) 23-33.

- [4] E. Egerváry, *On combinatorial properties of matrices*, Matematikai Lapok **38** (1931) 16-28.
- [5] O. Favaron, *Very well-covered graphs*, Discrete Mathematics **42** (1982) 177-187.
- [6] A. Finbow, B. Hartnell, R. J. Nowakowski, *A characterization of well-covered graphs of girth 5 or greater*, Journal of Combinatorial Theory Ser. B **57** (1993) 44-68.
- [7] D. König, *Graphen und Matrizen*, Matematikai Lapok **38** (1931) 116-119.
- [8] E. Korach, *On dual integrality, min-max equalities and algorithms in combinatorial programming*, University of Waterloo, Department of Combinatorics and Optimization, Ph.D. Thesis, 1982.
- [9] E. Korach, T. Nguyen, B. Peis, *Subgraph characterization of red/blue-split graphs and König-Egerváry graphs*, Proceedings of the Seventeenth Annual ACM-SIAM Symposium on Discrete Algorithms, ACM Press (2006) 842-850.
- [10] V. E. Levit, E. Mandrescu, *Well-covered and König-Egerváry graphs*, Congressus Numerantium **130** (1998) 209-218.
- [11] V. E. Levit, E. Mandrescu, *Combinatorial properties of the family of maximum stable sets of a graph*, Discrete Applied Mathematics **117** (2002) 149-161.
- [12] V. E. Levit, E. Mandrescu, *On α^+ -stable König-Egerváry graphs*, Discrete Mathematics **263** (2003) 179-190.
- [13] V.E. Levit, E. Mandrescu, *Square-stable and well-covered graphs*, Acta Universitatis Apulensis **10** (2005) 297-308.
- [14] V. E. Levit, E. Mandrescu, *On α -critical edges in König-Egerváry graphs*, Discrete Mathematics **306** (2006) 1684-1693.
- [15] V. E. Levit, E. Mandrescu, *Partial unimodality for independence polynomials of König-Egerváry graphs*, Congressus Numerantium **179** (2006) 109-119.

- [16] V. E. Levit, E. Mandrescu, *Triangle-free graphs with uniquely restricted maximum matchings and their corresponding greedoids*, Discrete Applied Mathematics **155** (2007) 2414-2425.
- [17] V. E. Levit, E. Mandrescu, *Critical independent sets and König-Egerváry graphs*, E-print: arXiv:0906.4609v2 [math.CO]
- [18] L. Lovász, *Ear decomposition of matching covered graphs*, Combinatorica **3** (1983) 105-117.
- [19] L. Lovász, M. D. Plummer, *Matching Theory*, Annals of Discrete Mathematics **29** (1986) North-Holland.
- [20] M. D. Plummer, *Some covering concepts in graphs*, Journal of Combinatorial Theory **8** (1970) 91-98.
- [21] E. Prisner, J. Topp, P. D. Vestergaard, *Well-covered simplicial, chordal, and circular arc graphs*, Journal of Graph Theory **21** (1996) 113-119.
- [22] B. Randerath, L. Volkman, *Simplicial graphs and relationships to different graph invariants*, Ars Combinatoria **46** (1997) 211-217.
- [23] G. Ravindra, *Well-covered graphs*, Journal of Combinatorial Information System Sciences **2** (1977) 20-21.
- [24] F. Sterboul, *A characterization of the graphs in which the transversal number equals the matching number*, Journal of Combinatorial Theory Ser. B **27** (1979) 228-229.

Vadim E. Levit
Department of Mathematics and Computer Science
Ariel University Center of Samaria
Ariel 40700, Israel
email:levitv@ariel.ac.il

Eugen Mandrescu
Department of Computer Science
Holon Institute of Technology
52 Golomb Str., Holon, Israel
email:eugen_m@hit.ac.il